

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 221/10, A61K 31/435, C07D 498/04, 471/04, 409/12, 401/12 // (C07D 498/04, 221:00, 265:00) (C07D 471/04, 241:00, 221:00)		A1	(11) International Publication Number: WO 97/40015 (43) International Publication Date: 30 October 1997 (30.10.97)
(21) International Application Number: PCT/US97/06676 (22) International Filing Date: 23 April 1997 (23.04.97) (30) Priority Data: 08/636,662 23 April 1996 (23.04.96) US (60) Parent Application or Grant (63) Related by Continuation US 08/636,662 (CIP) Filed on 23 April 1996 (23.04.96) (71) Applicant (for all designated States except US): NEUROGEN CORPORATION [US/US]; 35 Northeast Industrial Road, Branford, CT 06405 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): CHEN, Xi [US/US]; Apartment 72, 288 Old Foxon Road, New Haven, CT 06513 (US). YUAN, Jun [US/US]; 26 Cedar Road, Clinton, CT 06413 (US). THURKAUF, Andrew [US/US]; 16 Fox Den Road, Danbury, CT 06811 (US).		(74) Agent: SARUSSI, Steven, J.; McDonnell Boehnen Hulbert & Berghoff, 300 South Wacker Drive, Chicago, IL 60606 (US). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published With international search report.	

(54) Title: TRICYCLIC AMINOALKYL CARBOXAMIDES; NOVEL DOPAMINE D₃ RECEPTOR SUBTYPE SPECIFIC LIGANDS

(57) Abstract

Disclosed are compounds of formula (I) or the pharmaceutically acceptable acid addition salts thereof wherein: R₁, R₄ represent organic or inorganic groups; A represents an alkylene group of 2 to 6 carbon atoms optionally substituted with one or more alkyl groups having from 1



to 4 carbon atoms; Y represents methylene, oxygene, sulfur, or NH; and X is either N, C or CH, which compounds are useful in the treatment of affective disorders such as schizophrenia, depression, Alzheimer's disease, movement disorders such as Parkinsonism and dystonia, and other disorders which respond to dopaminergic blockade such as substance abuse and obsessive compulsive disorders. Further, compounds of this invention may be useful in treating the extrapyramidal side effects associated with the use of conventional neuroleptic agents.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

**Tricyclic aminoalkylcarboxamides;
Novel Dopamine D₃ Receptor Subtype Specific Ligands**

BACKGROUND OF THE INVENTION

5 **Field of the invention**

 This invention relates to tricyclic aminoalkylcarboxamide derivatives which selectively bind to brain dopamine receptor subtypes. It also relates to pharmaceutical compositions comprising such compounds. It further relates to the use of such compounds in the treatment or prevention of various neuropsychochological disorders such as schizophrenia and other central nervous system diseases.

10 **Description of the Related Art**

 The therapeutic effect of conventional antipsychotics, known as neuroleptics, is generally believed to be exerted through blockade of dopamine receptors. However, neuroleptics are frequently responsible for undesirable extrapyramidal side effects (EPS) and tardive dyskinesias, which are attributed to blockade of D₂ receptors in the striatal region of the brain. The dopamine D₃ receptor subtype has recently been identified (Sokoloff et al., Nature, 347, 146 (1990)). Its unique localization in limbic brain areas and its differential recognition of various antipsychotics suggest that the D₃ receptor plays a role in the etiology of schizophrenia. Selective D₃ antagonists thus are expected to be effective antipsychotics free from the neurological side effects displayed by conventional neuroleptics.

 U.S. Patent No. 5,395,835 discloses N-aminoalkyl-2-naphthalamides described to have affinity at dopamine D₃ receptors.

 Murray et al., Bioorg. Med. Chem. Let. 5: 219 (1995), describe 4-carboxamidobiphenyls said to have affinity at dopamine D₃ receptors.

 International Patent Application WO 9511903 discloses tricyclic diazaphenanthrene carboxamides having affinity for 5-HT_{1A} receptors.

 European Patent Application EP 206225 A2 discloses aryl alkyl azatricyclic compounds.


SUMMARY OF THE INVENTION

This invention provides novel compounds of Formula I which interact with dopamine receptor subtypes. Thus, the invention provides compounds of general Formula I useful in the treatment and/or prevention of various neuropsychological disorders. The invention also provides pharmaceutical compositions comprising compounds of Formula I.

Since dopamine D₃ receptors are concentrated in the limbic system (Taubes, Science, 265: 1034 (1994)) which controls cognition and emotion, compounds which interact with these receptors also have utility in the treatment of cognitive disorders. Such disorders include cognitive deficits which are a significant component of the negative symptoms (social withdrawal and unresponsiveness) of schizophrenia. Other disorders involving memory impairment or attention deficit disorders can also be treated with the compounds of this invention which interact specifically with the dopamine D₃ receptor subtype.

The invention thus encompasses the use of such compounds and compositions in the treatment of affective disorders such as schizophrenia, depression, Alzheimer's disease and certain movement disorders such as Parkinsonism and dystonia. Compounds of this invention are also useful in treating the extrapyramidal side effects associated with the use of conventional neuroleptic agents. Further, the compounds of the present invention are useful for the treatment of other disorders which respond to dopaminergic blockade such as substance abuse and obsessive compulsive disorder.

Accordingly, a broad embodiment of the invention is directed to a compound of Formula I:



I

or the pharmaceutically acceptable acid addition salts thereof wherein:

R₁, and R₂ are the same or different and represent hydrogen, C₁-C₆ alkyl, halogen, hydroxy, amino, cyano, nitro, trifluoromethyl, trifluoromethoxy, C₁-C₆ alkoxy, -O₂CR', -NHCOR', -COR', -SO_mR', where R' is C₁-C₆ alkyl and wherein m is 0, 1 or 2; or

R₁ and R₂ independently represent -CONR'R'', or -NR'R'' where R' and R'' independently represent hydrogen or C₁-C₆ alkyl;

R₃ is hydrogen or lower alkyl;

A represents an alkylene group of 2 to 6 carbon atoms optionally substituted with one or more alkyl groups having from 1 to 4 carbon atoms;

Y represents methylene, oxygen, sulfur, or NH;

5 X is either N, C, or CH; and

R₄ is biphenyl, naphthyl, 1 or 2 anthracenyl, 1 or 2 biphenylenyl, dibenzofuranyl, dibenzothiophenyl, fluorenyl, fluorenonyl, anthroquinonyl, 2-quinoxalinyll or azabiphenyl, each of which is optionally substituted independently by up to three groups selected from halogen, C₁-C₆ alkyl, C₁-C₄ alkoxy, alkylthio, hydroxy, amino, 10 monoalkylamino, dialkylamino, cyano, nitro, trifluoromethyl or trifluoromethoxy.


Thus, the invention provides compounds of Formula I useful in the treatment and/or prevention of neuropsychochological disorders including, but not limited to, schizophrenia, Alzheimer's disease, mania, dementia, depression, anxiety, compulsive behavior, substance 15 abuse, memory impairment, cognitive deficits, Parkinson-like motor disorders such as dystonia, and motion disorders and extrapyramidal side effects related to the use of neuroleptic agents.

DETAILED DESCRIPTION OF THE INVENTION

Compounds of the present invention demonstrate high affinity and selectivity in binding to the D₃ receptor subtype. They are therefore useful in treatment of schizophrenia, psychotic depression and mania. Other dopamine-mediated diseases such as Parkinsonism and tardive dyskinesias can also be treated directly or indirectly by modulation of D₃ receptors.

Furthermore, compounds of this invention can be used in treatment of memory-impairment or Alzheimer's disease by modulation of D₃ receptors which selectively exist in limbic area known to control emotion and cognitive functions. The compounds of the present invention are also useful for the treatment of other disorders which respond to dopaminergic blockade such as substance abuse (Caine and Koob, Science, 260: 1814 (1993)) and obsessive compulsive disorder (Goodman et al., Clin. Psychopharmacol., 7: 35 (1992)). The compounds of the invention interact with dopamine receptor subtypes resulting in the pharmacological activity of these compounds.

In addition to compounds of general Formula I described above, the invention encompasses compounds of Formula II:




II

wherein R₁-R₄, and A are as defined above for Formula I.

Preferred compounds of formula II include those where R₃ is hydrogen, methyl or ethyl, and A is alkylene of 3-5 carbon atoms. Particularly preferred compounds of Formula II are those where R₃ is hydrogen, and A is butylene.

The invention also provides compounds of Formula IIA:




IIA

wherein R₁-R₄, and A are as defined above for Formula I.

Preferred compounds of Formula IIA include those where R₃ is hydrogen, methyl or ethyl, and A is alkylene of 3-5 carbon atoms. Particularly preferred compounds of Formula IIA are those where R₃ is hydrogen, and A is butylene.

5

The invention also provides compounds of Formula IIB:




IIB

wherein R₁-R₄, and A are as defined above for Formula I.

10 Preferred compounds of Formula IIB include those where R₃ is hydrogen, methyl or ethyl, and A is alkylene of 3-5 carbon atoms. Particularly preferred compounds of Formula IIB are those where R₃ is hydrogen, and A is butylene.

The invention also provides compounds of Formula III:



III

wherein R₁-R₄, A and Y are as defined above for Formula I.

Preferred compounds of Formula III include those where R₃ is hydrogen, methyl or ethyl, and A is alkylene of 3-5 carbon atoms. More preferred compounds of Formula III are those where Y is oxygen or sulfur, R₃ is hydrogen, methyl or ethyl, and A is alkylene of 3-5 carbon atoms. Particularly preferred compounds of Formula III are those where Y is oxygen or sulfur, R₃ is hydrogen, and A is butylene.

20


In addition, the invention encompasses compounds of Formula IIIA, i.e., compounds of Formula III where Y is methylene.

In addition, the invention encompasses compounds of Formula IIIB, i.e., compounds of Formula III where Y is oxygen.

In addition, the invention encompasses compounds of Formula IIIC, i.e., compounds of Formula III where Y is sulfur.

5 In addition, the invention encompasses compounds of Formula IIID, i.e., compounds of Formula III where Y is NH.

Preferred compounds of the invention include R₄ groups selected from the following:



10

When a compound of the invention is obtained as a mixture of enantiomers, these enantiomers may be separated, when desired, by conventional methods such as crystallization in the presence of a resolving agent, or chromatography, for example using a chiral HPLC column.

15

Representative compounds of the present invention, which are encompassed by Formula I, include, but are not limited to the compounds in Table 1 and their pharmaceutically acceptable salts. The present invention also encompasses prodrugs, such as acylated prodrugs, of the compounds of Formula I. Those skilled in the art will recognize

various synthetic methodologies which may be employed to prepare non-toxic pharmaceutically acceptable addition salts and prodrugs of the compounds encompassed by Formula I.

Non-toxic pharmaceutically acceptable salts include salts of acids such as hydrochloric, phosphoric, hydrobromic, sulfuric, sulfinic, formic, toluenesulfonic, methanesulfonic, nitric, benzoic, citric, tartaric, maleic, hydroiodic, alkanolic such as acetic, $\text{HOOC}-(\text{CH}_2)_n-\text{COOH}$ where n is 0-4, and the like. Those skilled in the art will recognize a wide variety of non-toxic pharmaceutically acceptable addition salts.

By "alkyl" and "lower alkyl" is meant straight and branched chain alkyl groups having from 1-6 carbon atoms, e.g., C_1 - C_6 alkyl.

By "lower alkoxy" and "alkoxy" is meant straight and branched chain alkoxy groups having from 1-6 carbon atoms, e.g., C_1 - C_6 alkoxy.

By halogen is meant fluorine, chlorine, bromine and iodine.

Representative examples of compounds according to the invention are shown in Table 1 below. The number below each compound is its compound number. Each of these compounds may be prepared according to the general reaction Scheme I set forth below.

The invention also pertains to the use of compounds of general Formula I in the treatment of neuropsychological disorders. The pharmaceutical utility of compounds of this invention are indicated by the following assays for dopamine receptor subtype affinity.

ASSAY FOR D₂ AND D₃ RECEPTOR BINDING ACTIVITY

Pellets of COS cells containing recombinantly produced D₂ or D₃ receptors from African Green monkey were used for the assays. The sample is homogenized in 100 volumes (w/vol) of 0.05 M Tris HCl buffer at 4° C and pH 7.4. The sample is then centrifuged at 30,000 x g and resuspended and rehomogenized. The sample is then centrifuged as described and the final tissue sample is frozen until use. The tissue is resuspended 1:20 (wt/vol) in 0.05 M Tris HCl buffer containing 100 mM NaCl.

Incubations are carried out at 48°C and contain 0.4 ml of tissue sample, 0.5 nM ³H-YM 09151-2 and the compound of interest in a total incubation of 1.0 ml. Nonspecific binding is defined as that binding found in the presence of 1 mM spiperone; without further additions, nonspecific binding is less than 20% of total binding. The binding characteristics of representative compounds of the invention for D₂ and D₃ receptor subtypes are shown in Table 1 for rat striatal homogenates.

TABLE I

Compound Number ¹	D ₂ K _i (nM)	D ₃ K _i (nM)
1	127	8
6	165	17
7	377	5
8	250	12
13	362	10
14	198	9

¹ Compound numbers correspond to the compounds described below in the examples.

The binding characteristics of compounds of Formula I for the D₃ receptor, expressed in nM, generally range from about 0.5 nanomolar (nM) to about 50 nanomolar (nM). These compounds typically have binding constants for the D₂ receptor of from at least about 100 nM to more than 500 nM. Thus, the compounds of the invention are generally at least about 10, preferably at least about 20, and most preferably at least about 25 time more selective for the D₃ receptor than the D₂ receptor. Even more preferably, these compounds are at least 20, and more preferably at least 25-50, times more selective for the D₃ receptor than the D₂ receptor.

The compounds of general Formula I may be administered orally, topically, parenterally, by inhalation or spray or rectally in dosage unit formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intrasternal injection or infusion techniques. In addition, there is provided a pharmaceutical formulation comprising a compound of general Formula I and a pharmaceutically acceptable carrier. One or more compounds of general Formula I may be present in association with one or more non-toxic pharmaceutically acceptable carriers and/or diluents and/or adjuvants and if desired other active ingredients. The pharmaceutical compositions containing compounds of general Formula I may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsion, hard or soft capsules, or syrups or elixirs.

Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide pharmaceutically elegant and

palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients which are suitable for the manufacture of tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, corn starch, or alginic acid; binding agents, for example starch, gelatin or acacia, and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed.

Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin or olive oil.

Aqueous suspensions contain the active materials in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydropropyl methylcellulose; sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide, for example, lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives, for example ethyl, or n-propyl p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, and one or more sweetening agents, such as sucrose or saccharin.

Oily suspensions may be formulated by suspending the active ingredients in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and flavoring agents may be added to provide palatable oral preparations. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional
5 excipients, for example sweetening, flavoring and coloring agents, may also be present.

Pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally-occurring gums, for example gum acacia or gum tragacanth, naturally-occurring
10 phosphatides, for example soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol, anhydrides, for example sorbitan monoleate, and condensation products of the and partial esters with ethylene oxide, for example polyoxyethylene sorbitan monoleate. The emulsions may also contain sweetening and flavoring agents.

Syrups and elixirs may be formulated with sweetening agents, for example glycerol,
15 propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative and flavoring and coloring agents. The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleaginous suspension. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may
20 also be sterile injectable solution or suspension in a non-toxic parentally acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including
25 synthetic mono or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

The compounds of general Formula I may also be administered in the form of suppositories for rectal administration of the drug. These compositions can be prepared by
30 mixing the drug with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials are cocoa butter and polyethylene glycols.

Compounds of general Formula I may be administered parenterally in a sterile medium. The drug, depending on the vehicle and concentration used, can either be suspended

or dissolved in the vehicle. Advantageously, adjuvants such as local anesthetics, preservatives and buffering agents can be dissolved in the vehicle.


Dosage levels of the order of from about 0.1 mg to about 140 mg per kilogram of body weight per day are useful in the treatment of the above-indicated conditions (about 0.5 mg to about 7 g per patient per day). The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. Dosage unit forms will generally contain between from about 1 mg to about 500 mg of an active ingredient.


It will be understood, however, that the specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, route of administration, and rate of excretion, drug combination and the severity of the particular disease undergoing therapy.

15 Preparation of tricyclic aminoalkylcarboxamides

The compounds of Formula I, and the pharmaceutically acceptable acid addition salts thereof, may be prepared according to the reactions shown below in Scheme 1.

Scheme 1





wherein R₁-R₄, A, X and Y are as defined above for Formula I.

As shown, an N-aminoalkyl tricyclic amine of structure 17 may be prepared by the
 5 condensation of an appropriate tricyclic amine derivative (16) with an N-haloalkylphthalimide. This initial reaction, which is followed by treatment with hydrazine hydrate, may be carried out at room temperature in the presence of a base in a solvent such as acetonitrile. The resulting compound of structure 17 may be condensed in the presence of
 10 base with an activated acid derivative, such as an acid chloride (R₄COCl) or anhydride to provide secondary carboxamides of structure 18. When it is desirable that R₃ be other than hydrogen, i.e., lower alkyl, a compound of formula 18 may be alkylated with an alkyl halide in the presence of base.

Where they are not commercially available, the haloalkylphthalimides may be prepared by literature procedures or procedures analogous to those described in literature.
 15 The 1,2,3,4-tetrahydrobenz[f]isoquinolines of general structure 16 used in preparing compounds of Formula IIA can be prepared according to the method of Kessar et al. (Indian J. Chem., 12(2), 113-116, 1974). The 1,2,3,4,5,6-hexahydrobenz[f]isoquinolines of general structure 16 used in preparing compounds of general Formula IIB can be prepared according to the method of Menard et al. (Can. J. Chem., 52, 2316-26, 1974). The bridged 4-
 20 phenylpiperazines of general structure 16 used in preparing compounds of general Formula III can be prepared according to the methods of Baxter and Richards (J. Med. Chem., 15, 351-357, 1972), Baxter and Reitz (International Application Publication No. WO 95/11903) or Fried and Potoski (U.S. Patent No. 4,089,958).

Those having skill in the art will recognize that the starting materials may be varied
 25 and additional steps employed to produce compounds encompassed by the present invention.

as demonstrated by the following examples. In some cases protection of certain reactive functionalities may be necessary to achieve some of the above transformations. In general the need for such protecting groups will be apparent to those skilled in the art of organic synthesis as well as the conditions necessary to attach and remove such groups.

5 The disclosures in this application of all articles and references, including patents, are incorporated herein by reference.


The invention is illustrated further by the following examples which are not to be construed as limiting the invention in scope or spirit to the specific procedures described in them. These examples illustrate the presently preferred methods for preparing the compounds
10 of the invention.

Example 1

1. 2-(4-aminobutyl)-1,2,3,4-tetrahydrobenz[f]isoquinoline

A solution of 6 g (30.6 mmol) of 1,2,3,4-tetrahydrobenz[f]isoquinoline and 8.70 g (31
15 mmol) of N-(4-bromobutyl)phthalimide in 100 ml of acetonitrile containing 5 g of potassium carbonate was heated at reflux for 6 h. Upon cooling the reaction mixture was washed with water, dried and concentrated. The resulting material was dissolved in 60 ml of 95% ethanol containing 15 ml of hydrazine hydrate and heated to reflux for 4h. Upon cooling the reaction mixture was concentrated and the resulting solid was stirred for 30 min with diethyl ether,
20 filtered and the filtrate concentrated to provide 7.1 g (86%) of the desired N-aminoalkyl tricyclic amine.

2. N-{4-(1,2,3,4-tetrahydrobenz[f]isoquinolin-2-yl)butyl}-quinoline-3-carboxamide hydrochloride



Compound 1

To a solution of 130 mg of 2-(4-aminobutyl)-1,2,3,4-tetrahydrobenz[f]isoquinoline and triethylamine (0.3 ml) in 5 ml of pentene stabilized chloroform was added in one portion to a solution 93 mg of quinoline-3-carbonylchloride in 5 ml of pentene stabilized chloroform.
30 The resulting mixture was washed with 2N NaOH, dried (Na₂SO₄) and concentrated. The resulting mixture was purified by chromatography on silica gel eluting with 10% methanol in


chloroform to provide the desired product (131 mg, 54 %). The hydrochloride salt was prepared in isopropanol.

Example 2

5 The following compounds are prepared essentially according to the procedures set forth above. The structure for each of the following compounds appears immediately below the corresponding compound name.


(a) N-{4-(1,2,3,4-tetrahydrobenz[f]isoquinolin-2-yl)butyl}-anthracene-2-carboxamide hydrochloride (Compound 2)

10




(b) N-{4-(1,2,3,4-tetrahydrobenz[f]isoquinolin-2-yl)butyl}-biphenylene-2-carboxamide hydrochloride (Compound 3)


15




(c) N-{4-(1,2,3,4-tetrahydrobenz[f]isoquinolin-2-yl)butyl}-fluorene-3-carboxamide hydrochloride (Compound 4)




(d) N-{4-(1,2,3,4-tetrahydrobenz[f]isoquinolin-2-yl)butyl}-naphthalene-2-carboxamide hydrochloride (Compound 5)




5 (e) N-{4-(1,2,3,4-tetrahydrobenz[f]isoquinolin-2-yl)butyl}-fluoren-9-one-3-carboxamide hydrochloride (Compound 6)




(f) N-{4-(1,2,3,4-tetrahydrobenz[f]isoquinolin-2-yl)butyl}-dibenzothiophene-2-carboxamide hydrochloride (Compound 7)




(g) N-{4-(1,2,3,4-tetrahydrobenz[f]isoquinolin-2-yl)butyl}-quinoxaline-2-carboxamide hydrochloride (Compound 8)




(h) N-{4-(1,2,3,4-tetrahydrobenz[f]isoquinolin-2-yl)butyl}-anthraquinone-2-carboxamide hydrochloride (Compound 9)




5 (i) N-{4-(1,2,3,4-tetrahydrobenz[f]isoquinolin-2-yl)butyl}-biphenyl-4-carboxamide hydrochloride (Compound 10)




10 (j) 2-[N-{4-(1,2,3,4-tetrahydrobenz[f]isoquinolin-2-yl)butyl}-4-carboxamidophenyl]pyridine hydrochloride (Compound 11)




(k) N-{4-(1,2,3,4,4a,5-hexahydropyrazino[2,1-c][1,4]benzoxazin-2-yl)butyl}-fluorene-2-carboxamide hydrochloride (Compound 12)



(l) N-{4-(1,2,3,4,5,6-hexahydrobenz[f]isoquinolin-2-yl)butyl}-biphenyl-4-carboxamide hydrochloride (Compound 13)




5 (m) N-{4-(2,3,4,4a,5,6-hexahydro-1H-pyrazino[1,2-a]quinolin-2-yl)butyl}-fluorene-2-carboxamide hydrochloride (Compound 14)



The invention and the manner and process of making and using it, are now described
10 in such full, clear, concise and exact terms as to enable any person skilled in the art to which it
pertains, to make and use the same. It is to be understood that the foregoing describes
preferred embodiments of the present invention and that modifications may be made therein
without departing from the spirit or scope of the present invention as set forth in the claims.
To particularly point out and distinctly claim the subject matter regarded as invention, the
15 following claims conclude this specification.

What is claimed is:

1. A compound of the formula:



or the pharmaceutically acceptable acid addition salts thereof wherein:

- 5 R_1 , and R_2 are the same or different and represent hydrogen, C_1 - C_6 alkyl, halogen, hydroxy, amino, cyano, nitro, trifluoromethyl, trifluoromethoxy, C_1 - C_6 alkoxy, $-O_2CR'$, $-NHCOR'$, $-COR'$, $-SO_mR'$, where R' is C_1 - C_6 alkyl and wherein m is 0, 1 or 2; or R_1 and R_2 independently represent $-CONR'R''$, or $-NR'R''$ where R' and R'' independently represent hydrogen or C_1 - C_6 alkyl;

- 10 R_3 is hydrogen or lower alkyl;

A represents an alkylene group of 2 to 6 carbon atoms optionally substituted with one or more alkyl groups having from 1 to 4 carbon atoms;


Y represents methylene, oxygen, sulfur, or NH;

X is N, C or CH; and

- 15 R_4 is biphenyl, naphthyl, 1 or 2 anthracenyl, 1 or 2 biphenylenyl, dibenzofuranyl, dibenzothiophenyl, fluorenyl, fluorenonyl, anthroquinonyl, 2-quinoxaliny or azabiphenyl, each of which is optionally substituted independently by up to three groups selected from halogen, C_1 - C_6 alkyl, C_1 - C_4 alkoxy, alkylthio, hydroxy, amino, monoalkylamino, dialkylamino, cyano, nitro, trifluoromethyl or trifluoromethoxy.

20


2. A compound according to Claim 1, which has the formula:



3. A compound according to Claim 2, wherein R_3 is hydrogen, methyl or ethyl;

- 25 and A is alkylene of 3-5 carbon atoms.

4. A compound of the formula:




wherein

- 5 R₁, and R₂ are the same or different and represent hydrogen, C₁-C₆ alkyl, halogen, hydroxy, amino, cyano, nitro, trifluoromethyl, trifluoromethoxy, C₁-C₆ alkoxy, -O₂CR', -NHCOR', -COR', -SO_mR', where R' is C₁-C₆ alkyl and wherein m is 0, 1 or 2; or R₁ and R₂ independently represent -CONR'R'', or -NR'R'' where R' and R'' independently represent hydrogen or C₁-C₆ alkyl;
- 10 R₃ is hydrogen or lower alkyl;
- A represents an alkylene group of 2 to 6 carbon atoms optionally substituted with one or more alkyl groups having from 1 to 4 carbon atoms;
- Y represents methylene, oxygen, sulfur, or NH; and
- R₄ is biphenyl, naphthyl, 1 or 2 anthracenyl, 1 or 2 biphenylenyl, dibenzofuranyl, 15 dibenzothiophenyl, fluorenyl, fluorenonyl, anthroquinonyl, 2-quinoxalinyll or azabiphenyl, each of which is optionally substituted independently by up to three groups selected from halogen, C₁-C₆ alkyl, C₁-C₄ alkoxy, alkylthio, hydroxy, amino, monoalkylamino, dialkylamino, cyano, nitro, trifluoromethyl or trifluoromethoxy.
- 20 5. A compound according to Claim 4, wherein R₃ is hydrogen and A is alkylene of 3-5 carbon atoms.


6. A compound according to Claim 4, wherein A is butylene.

7. A compound according to Claim 1, which has the formula



8. A compound according to Claim 7, wherein R₃ is hydrogen and A is alkylene
5 of 3-5 carbon atoms.

9. A compound of the formula:



where

- 10 R₁, and R₂ are the same or different and represent hydrogen, C₁-C₆ alkyl, halogen, hydroxy, amino, cyano, nitro, trifluoromethyl, trifluoromethoxy, C₁-C₆ alkoxy, -O₂CR', -NHCOR', -COR', -SO_mR', where R' is C₁-C₆ alkyl and wherein m is 0, 1 or 2; or R₁ and R₂ independently represent -CONR'R'', or -NR'R'' where R' and R'' independently represent hydrogen or C₁-C₆ alkyl;

- 15 R₃ is hydrogen or lower alkyl;

A represents an alkylene group of 2 to 6 carbon atoms optionally substituted with one or more alkyl groups having from 1 to 4 carbon atoms;

Y represents methylene, oxygen, sulfur, or NH;

- 20 R₄ is biphenyl, naphthyl, 1 or 2 anthracenyl, 1 or 2 biphenylenyl, dibenzofuranyl, dibenzothiophenyl, fluorenyl, fluorenonyl, anthroquinonyl, 2-quinoxaliny or azabiphenyl, each of which is optionally substituted independently by up to three groups selected from halogen, C₁-C₆ alkyl, C₁-C₄ alkoxy, alkylthio, hydroxy, amino, monoalkylamino, dialkylamino, cyano, nitro, trifluoromethyl or trifluoromethoxy.

10. A compound according to Claim 7, wherein R₃ is hydrogen, and A is alkylene of 3-5 carbon atoms.

11. A compound according to Claim 10, wherein A is butylene.

5

12. A compound according to Claim 1, which is N-{4-(1,2,3,4-tetrahydrobenz[f]isoquinolin-2-yl)butyl}-quinoline-3-carboxamide.

13. A compound according to Claim 1, which is N-{4-(1,2,3,4-tetrahydrobenz[f]isoquinolin-2-yl)butyl}-anthracene-2-carboxamide.

10

14. A compound according to Claim 1, which is N-{4-(1,2,3,4-tetrahydrobenz[f]isoquinolin-2-yl)butyl}-biphenylene-2-carboxamide.

15

15. A compound according to Claim 1, which is N-{4-(1,2,3,4-tetrahydrobenz[f]isoquinolin-2-yl)butyl}-biphenylene-2-carboxamide.

16. A compound according to Claim 1, which is N-{4-(1,2,3,4-tetrahydrobenz[f]isoquinolin-2-yl)butyl}-fluorene-3-carboxamide.

20

17. A compound according to Claim 1, which is N-{4-(1,2,3,4-tetrahydrobenz[f]isoquinolin-2-yl)butyl}-naphthalene-2-carboxamide hydrochloride.

18. A compound according to Claim 1, which is N-{4-(1,2,3,4-tetrahydrobenz[f]isoquinolin-2-yl)butyl}-fluoren-9-one-3-carboxamide hydrochloride.

25

19. A compound according to Claim 1, which is N-{4-(1,2,3,4-tetrahydrobenz[f]isoquinolin-2-yl)butyl}-dibenzothiophene-2-carboxamide hydrochloride.

20. A compound according to Claim 1, which is N-{4-(1,2,3,4-tetrahydrobenz[f]isoquinolin-2-yl)butyl}-quinoxaline-2-carboxamide hydrochloride.

30

21. A compound according to Claim 1, which is N-{4-(1,2,3,4-tetrahydrobenz[f]isoquinolin-2-yl)butyl}-anthraquinone-2-carboxamide hydrochloride.

22. A compound according to Claim 1, which is N-{4-(1,2,3,4-tetrahydrobenz[f]isoquinolin-2-yl)butyl}-biphenyl-4-carboxamide hydrochloride.

5 23. A compound according to Claim 1, which is 2-[N-{4-(1,2,3,4-tetrahydrobenz[f]isoquinolin-2-yl)butyl}-4-carboxamidophenyl]pyridine hydrochloride.

24. A compound according to Claim 1, which is N-{4-(1,2,3,4,5,6-hexahydrobenz[f]isoquinolin-2-yl)butyl}-biphenyl-4-carboxamide hydrochloride.

10

25. A compound according to Claim 1, which is N-{4-(1,2,3,4,4a,5-hexahydropyrazino[2,1-c][1,4]benzoxazin-2-yl)butyl}-fluorene-2-carboxamide hydrochloride.

15 26. A compound according to Claim 1, which is N-{4-(2,3,4,4a,5,6-hexahydro-1H-pyrazino[1,2-a]quinolin-2-yl)butyl}-fluorene-2-carboxamide hydrochloride.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 97/06676

A. CLASSIFICATION OF SUBJECT MATTER IPC 6 C07D221/10 A61K31/435 C07D498/04 C07D471/04 C07D409/12 C07D401/12 //(C07D498/04,265:00,221:00),(C07D471/04,241:00, 221:00) According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 6 C07D Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 95 11903 A (ORTHO PHARMACEUTICAL CORPORATION) 4 May 1995 cited in the application see claims ---	1-4
A	WO 95 14006 A (THE BOARD OF GOVERNORS FOR HIGHER EDUCATION STATE OF RHODE ISLAND) 26 May 1995 see claims ---	1-4
A	US 3 647 791 A (ALBERTO ROSSI ET AL) 7 March 1972 see the whole document --- -/-	1-4
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "&" document member of the same patent family		
Date of the actual completion of the international search 25 July 1997		Date of mailing of the international search report 04.08.97
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl, Fax (+ 31-70) 340-3016		Authorized officer Henry, J

INTERNATIONAL SEARCH REPORT

Intern. Application No
PCT/US 97/06676

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	CHEMICAL ABSTRACTS, vol. 122, no. 19, 8 May 1995 Columbus, Ohio, US; abstract no. 239643m, page 1055; XP002036052 see abstract	1-4
A	& BIOORG.MED.CHEM.LETT., vol. 5, no. 3, 1995, pages 219-222, JOHN P. MURRAY ET AL: "A novel series of arylpiperazines with high affinity and selectivity for the dopamine D3 receptor." cited in the application ---	1-4
P,X	WD 96 23789 A (SANKYO COMPANY LIMITED) 8 August 1996 see abstract see page 29 - page 52 -----	1,9,25, 26

INTERNATIONAL SEARCH REPORT
Information on patent family members

Intern. Application No
PCT/US 97/06676

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9511903 A	04-05-95	US 5512566 A AU 7975794 A EP 0746556 A ZA 9408354 A	30-04-96 22-05-95 11-12-96 24-04-96
WO 9514006 A	26-05-95	AU 5615394 A EP 0729458 A	06-06-95 04-09-96
US 3647791 A	07-03-72	NONE	
WO 9623789 A	08-08-96	AU 4547796 A JP 8269057 A	21-08-96 15-10-96

THIS PAGE BLANK (USPTO)